







Liver Forum 5 November 10, 2016

Cysteamine Bitartrate Delayed-Release for the Treatment of Nonalcoholic Fatty Liver Disease in Children

Jeffrey B. Schwimmer, M.D.

Professor of Pediatrics, UC San Diego
Director, Fatty Liver Clinic, Rady Children's Hospital





DECISIONS

Drug Choice

Dose Choice



End Point

































CyNCh

End Point

Ist trial in pediatric liver disease designed with liver histology as the primary outcome





PRIMARY OUTCOME

End Point

The proportion of children with histologic improvement in NAFLD between the baseline liver biopsy and follow-up biopsy after 52 weeks of treatment, where improvement was defined as:

- I. decrease in NAS of 2 points or more and
- no worsening of fibrosis.



Who to Treat

HISTOLOGIC ENTRY CRITERIA

• NAS ≥ 4






Who to Treat

HISTOLOGIC ENTRY CRITERIA

• NAS ≥ 4







Dose Choice

WEIGHT BASED DOSING STRATA

Weight	Dose
≤ 65 kg	300 mg BID
>65 to 80 kg	375 mg BID
> 80 kg	450 mg BID






Who to Treat


End Point







CYSTEAMINE BITARTRATE DID NOT IMPROVE NAFLD ACTIVITY SCORE









CYSTEAMINE BITARTRATE IMPROVED ALT AND AST



Cysteamine Bitartrate






P < 0.01

P < 0.001








P < 0.01






CHILDREN < 65 KG RECEIVED SIGNIFICANTLY HIGHER DOSE (MG/KG)



P < 0.0001



HISTOLOGIC IMPROVEMENT BY DOSE (MG/KG)TERTILES







WEIGHT BASED DOSING STRATA

Weight	Dose	Capsules per day
≤ 65 kg	300 mg BID	8
>65 to 80 kg	375 mg BID	10
> 80 kg	450 mg BID	12








WEIGHT GAIN

- Lifestyle counseling c/w AAP Guidelines
- Mean weight gain = 7.1 kg
- 1.3 pounds per month




HISTOLOGIC IMPROVEMENT BY AGE







IMPROVEMENT IN TYPE 2 OR BORDERLINE PORTAL NASH WITH CYSTEAMINE BITARTRATE







QUESTIONS

Drug Choice

Dose Choice Who to Treat

End Point

